

## Quantification of nutrient content in above-ground biomass of young *Acacia mearnsii* De Wild., provenance Bodalla

Marcos Vinícius Winckler Caldeira<sup>a</sup>, Mauro Valdir Schumacher<sup>a</sup> and Peter Spathelf<sup>b\*</sup>

<sup>a</sup> Universidade Federal de Santa Maria – Departamento de Ciências Florestais, Av. Roraima S/N, Camobi, CEP: 97105-900, Santa Maria, RGS/Brazil  
<sup>b</sup> Forstdirektion Tübingen, Im Schloss, 72074 Tübingen, Germany

(Received 28 February 2001; accepted 28 November 2001)

**Abstract** – The present study deals with the quantification of the nutrient content of the Australian acacia (*Acacia mearnsii* De Wild.), provenance Bodalla, at the age of 28 months. The experiment is located at a site of low fertility of the AGROSETA company in Butiá, Rio Grande do Sul (Brazil). Above-ground biomass components (leaves, living branches, dead branches, bark and wood) of nine sampled trees were analysed. The nutrient contents in the total biomass of Bodalla were 182 kg ha<sup>-1</sup> of N, 8.17 of P, 104 of K, 66.7 of Ca, 16.2 of Mg, and 10.1 of S. In total, 42.6% of the dry matter accounted for leaves and living and dead branches, which in turn account for 74% of N, 72% of P, 63% of K, 68% of Ca, 69% of Mg, and 74% of S of the above-ground biomass. In the trunk (bark and wood), which represents the remaining 57.4% of the total above-ground biomass, 26% of N, 28% of P, 37% of K, 32% of Ca, 31% of Mg, and 26% of S were accumulated.

*Acacia mearnsii* De Wild. / biomass / nutrients / mineralmass / biogeochemical cycles

**Résumé** – Quantification des nutriments dans la biomasse aérienne de jeunes *Acacia mearnsii* De Wild., provenance Bodalla. Cette étude concerne la quantification, à l'âge de 28 mois, des nutriments dans la biomasse aérienne de l'Acacia australien (*Acacia mearnsii* De Wild.), provenance Bodalla. L'expérience est localisée dans un site de faible fertilité appartenant à la compagnie AGROSETA, à Butiá, Rio Grande do Sul (Brésil). Les compartiments de la biomasse aérienne (feuilles, branches vivantes, branches mortes, écorce et bois) de 9 arbres échantillons ont été analysés. Les quantités de nutriments dans la biomasse totale de Bodalla étaient de 182 kg ha<sup>-1</sup> de N, 8,17 de P, 104 de K, 66,7 de Ca, 16,2 de Mg et 10,1 de S. Au total, 42,6 % de la matière sèche était composée de feuilles et de branches vivantes et mortes, représentant 74 % de N, 72 % de P, 63 % de K, 68 % de Ca, 69 % de Mg, et 74 % de S par rapport à la biomasse aérienne totale. Dans le tronc (écorce et bois), qui représente les 57,4 % restant de la biomasse aérienne totale, se sont accumulés 26 % de N, 28 % de P, 37 % de K, 32 % de Ca, 31 % de Mg et 26 % de S.

*Acacia mearnsii* De Wild. / biomasse / nutriments / minéralomasse / cycles biogéochimiques

### 1. INTRODUCTION

Nutrient content in the above-ground biomass increases from boreal to tropical forests [20]. On the other hand, the amount of nutrients accumulated in litter and other above-ground deposits increases from tropical to boreal forests. This is generally due to the low activity of decomposing organisms in boreal forests inhibited by low temperatures and/or drought [14]. Furthermore, nutrient deficiency and water stress are seen to be a major constraint in forestry in the seasonal tropics [20], such as the Brazilian *Cerrado*.

In general, the annual absorption of nutrients in forests is lower than that of agricultural areas, but in forests most of the absorbed nutrients are returned to the site and only small quantities are retained in the forest biomass. According to Young and Carpenter [38], nutrient absorption is influenced by species, age, crown covering and edafo-climatic conditions. Furthermore, nutrient absorption in forest plantations is closely associated with the increase in biomass and attains its maximum in the initial stage of a rotation period [18]. After that stage, nutrient demand can generally be satisfied by translocation or internal cycling. Bellote et al. [6] showed that most of the biomass and nutrients accumulated by

\* Correspondence and reprints  
Tel.: +49 7071 602 213; fax: +49 7071 602 601; e-mail: Peter.Spathelf@forst.bwl.de

planted *Eucalyptus grandis* occurred between 2 and 5 years of age. When leaf area is expanding, nutrient uptake from the soil is high. After stand crown closure, nutrient allocation shifts from the crown (leaves) to the trunk [15, 20].

Nutrient export from a forest site is proportional to the quantity of biomass removed. Further nutrient losses can be due to erosion and outwashing after tree felling when the soil remains bare [20].

Total nutrient cycle in a forest ecosystem is represented by the sum of nutrients in the different ecosystem components such as above- and below-ground tree biomass, understorey vegetation, litter and soil [25]. In boreal forests, most of the nutrients can be found in the active tree tissues, such as leaves [37]. According to Van Den Driessche [34], conifers tend to have a higher proportion of leaf biomass than broad-leaved trees. In contrast to broad-leaved trees, a major percentage on total nutrient content can be found in the leaves of conifers, although nutrient concentration in the leaves of conifers is lower than in broad-leaved trees.

It seems to be a general observation that nutrient contents in tree compartments vary with species. Lugo [17] found significant differences in biomass and nutrient accumulation for N, P, and K in different tropical plantation species under similar edafo-climatic conditions in Costa Rica, thus emphasizing the different nutrient-use-efficiencies of the species involved.

Plantation species of the genus *Eucalyptus*, *Pinus*, or *Acacia* play a major role in Brazilian wood industry supply. Although industrial plantations only encompass  $7 \times 10^6$  ha (out of the  $553 \times 10^6$  ha of total forest area in Brazil,  $546 \times 10^6$  ha are natural forests), they contribute 60% to wood production [20]. *Acacia mearnsii* is considered to be an important tree species in Rio Grande do Sul, the southernmost state of Brazil. The total area with this species is approximately  $160 \times 10^3$  ha, planted mostly by small landowners.

The knowledge of nutrient quantity in the nutrient stock of the soil, above and below-ground biomass is of fundamental importance to the understanding of a forest ecosystem. A deeper insight into nutrient dynamics is also a precondition for guaranteeing ecological sustainability in these forest plantations [29]. In studies with *Eucalyptus* sp. it was shown that short rotations led to higher nutrient export per area per year than long rotations [26]. Poggiani [24] also calculated the length of time before the nutrient capital in the soil is exhausted in relation to nutrient quantities accumulated in forest biomass.

In the face of the lack of knowledge concerning nutrition and nutrient cycling of *Acacia mearnsii* [7] in Rio Grande do Sul, the objective of the present study is to quantify nutrient content of above-ground biomass in a 28-month-old stand of Australian acacia (*Acacia mearnsii* De Wild.), provenance Bodalla, as the first step in the analysis of *Acacia* plantations over a complete rotation period.

## 2. MATERIALS AND METHODS

### 2.1. Experimental area

The present study was carried out on an experimental site belonging to AGROSETA Florestal Company, which is located in the city of Butiá, Rio Grande do Sul (Brazil). The study site is situated in the so-called “Serra do Sudeste” (southeastern mountain range) of Rio Grande do Sul, with the following geographical coordinates: latitude  $30^{\circ} 07' 12''$  South and longitude  $51^{\circ} 57' 45''$  West. The study site is situated at an altitude of 40 m a.s.l. and the relief is slightly undulating.

According to Köppen, the climate of the region is Cfa type, i.e., subtropical humid [19]. Mean annual air temperature ranges from 18–19 °C, with mean annual temperatures of 24 °C in the hottest month (January) and 14 °C in the coldest month (July). Annual, January, and July precipitation are 1400 mm, 130 mm, and 120 mm, respectively.

The geological parent material of the study region is granite. The weathered granite substrate yields significant contents of clay. According to EMBRAPA [11], the soils of the study region belong to the soil mapping unit of “São Jerônimo” with high amounts of silicates. In general these dark-red podzolic soils are deep and have good drainage. Furthermore, they are acid and have small amounts of soil organic matter [11, 21]. According to the International Soil Taxonomy this soil can be classified as Ultisol.

### 2.2. Chemical analysis of soils

Low values of soil pH and poor soil organic matter content are typical of many sites in southern Brazil with plantations of *Acacia*. Another characteristic is the low content of soil available P which, in general, is due to the high P-fixing capacity of the soil [11, 21]. Al concentration (in the form of hydrous oxides) is high in these soils and correlates positively with a low cation exchange capacity [7]. Furthermore, the base saturation value below 50% is linked to the rather low fertility of the experimental site (*table I*). Chemical analysis of the sampling site was carried out using standard extraction methods at the Soil Department laboratory of the Federal University of Rio Grande do Sul in Porto Alegre (Brazil).

### 2.3. Provenance

The *Acacia* variety under study was Bodalla from New South Wales, Australia, with the following geographical coordinates: latitude  $36^{\circ} 11'$  South; longitude  $149^{\circ} 58'$  East (altitude: 15 m a.s.l.).

### 2.4. Determination of the nutrient concentration in above-ground biomass

The experimental stand was planted in August 1994 with an initial spacing of 1.7 m  $\times$  3.0 m. A homogeneous area was selected for this experiment according to the criteria (e.g., soil type, soil bulk density, and penetration resistance) of the soil mapping unit “São Jerônimo”. Four rectangular sampling areas of 18 m  $\times$  24 m (altogether 1728 m<sup>2</sup>) were established. Diameter distribution of the experimental stand is shown in *table II*. All diameters at breast height (dbh, in cm) were measured within the experimental area. The heights of 10% of the trees were measured. Also, a height (h, in m) estimation was carried out using the following model:  $\log h = (b_0 + b_1/dbh)^2 + 1.30$ .

**Table I.** Chemical analysis of the soil from the sampling site of *Acacia mearnsii*, provenance Bodalla (AGROSETA Florestal Company, experimental sites, Rio Grande do Sul, Brazil).

Soil horizon (depth) (cm)	pH (H <sub>2</sub> O) 1:1	soil organic matter (mg g <sup>-1</sup> )	avail. P (mg L <sup>-1</sup> )	avail. K (mg L <sup>-1</sup> )	exchang. Al <sup>3+</sup> (cmol(+) kg <sup>-1</sup> )
0–10	4.7	32	5.3	111	0.6
10–20	4.5	21	2.6	71.5	1.7
20–30	4.5	16	2.0	68.5	2.5
Soil horizon (depth) (cm)	exchang. Ca <sup>2+</sup> (cmol(+) kg <sup>-1</sup> )	exchang. Mg <sup>2+</sup> (cmol(+) kg <sup>-1</sup> )	CEC (cmol(+) kg <sup>-1</sup> )	Base saturation (%)	
0–10	2.2	1.5	4.6	44	
10–20	0.9	0.7	3.4	22	
20–30	0.5	0.4	3.5	12	

mg L<sup>-1</sup> = milligram per liter; CEC = cation exchange capacity.

**Table II.** Distribution of diameter at breast height in the experimental stand of *Acacia mearnsii*.

Diameter class (cm)								
3.0–4.0	4.1–5.0	5.1–6.0	6.1–7.0	7.1–8.0	8.1–9.0	9.1–10.0	10.1–11.0	11.1–12.0
6	21	55	51	63	37	11	2	3

Tree volume ( $v$ , in m<sup>3</sup>) was calculated with the equation:  $v = b_0 + b_1 \times dbh^2 \times h$ .

Thus, mean diameter and mean height, tree number, basal area, as well as tree volume over bark was calculated for each sampling area.

Nine trees (one tree in each diameter class) were sampled for subsequent nutrient analysis. In the intermediate part of the tree crown, leaves were collected at four cardinal points. Branches were separated from the trunk and classified as alive (green) and dead (dry). All the leaves were collected from living branches. Then, total fresh weight of leaves, living and dead branches, bark and trunk wood of the sampled trees were determined in the field.

The samples were dried in a circulation oven for 72 hours at temperatures ranging from 70 to 75 °C, until a constant weight was attained. Finally, samples were weighed with an analytical balance in order to obtain dry weight (dw). A disk 5 cm thick was removed at the mid-point of each trunk, as proposed by Young & Carpenter [38]. For each disk, the bark was separated from the wood. This measurement served as a reference for dw determination. The bark and wood samples were ground in a Wiley mill and then passed through a 1.0 mm sieve.

Biomass ( $Y$ ) of the trees was determined by using the following regression equations:

$$\ln Y = a + b \times \ln dbh \text{ (for leaves/ living branches);}$$

$$\text{and } \ln Y = a + b \times \ln dbh + c \times \ln h \text{ (for bark and wood) [7].}$$

$Y$  is expressed in Mg ha<sup>-1</sup> and  $dbh$  in cm;  $a$ ,  $b$ , and  $c$  are regression coefficients.

The nutrient concentrations of total N, P, K, Ca, Mg, and S in biomass were obtained using the methods of Tedesco et al. [32]. This method allows that the five macroelements can be determined with one solution of H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub>. The nutrients are restored, similar to the Kjeldahl method for N and with a nitric-perchloric solution for the other elements.

## 2.5. Nutrient quantification in above-ground biomass

Macronutrient stock (kg ha<sup>-1</sup>) in the above-ground biomass was calculated on the basis of the biomass estimation (kg ha<sup>-1</sup>) made by Caldeira [7] and the macronutrient concentrations (g kg<sup>-1</sup>) obtained in the present study. The sum of the values for each component provided the total nutrient content (kg ha<sup>-1</sup>) of above-ground biomass.

## 3. RESULTS AND DISCUSSION

Different nutrient concentrations in different tree species (or provenances) can be due to environmental conditions or genetic characteristics of the species [15, 20]. Nutrient concentrations of the different tree components are related to the production of above and below-ground biomass, stand density, and soil. The concentrations of N, P, K, Ca, Mg, and S in the components of above-ground biomass of Bodalla are shown in *table III*.

It is evident that most of the nutrients are concentrated in the leaves. Similar results were found by Tandon et al. [31] in Australian plantations of *Eucalyptus grandis*, Baggio [3] in *Mimosa scabrella*, Vezzani [35] in pure and mixed stands of *Eucalyptus saligna* and *Acacia mearnsii*, Pereira et al. [22] in *Acacia mearnsii*, and by George and Varghese [12] in *Eucalyptus globulus*. Nutrient concentration in leaves is influenced by diverse factors such as site conditions, age, position of leaves and season [34]. Bellote [5], who worked with *Eucalyptus grandis* in Brazil, observed that nutrient concentration in leaves varies with stand age and the season.

The high amount of N in the leaves of *Acacia mearnsii* is due to the capacity of this species to fix N. In tropical soils *Acacia mearnsii* is able to fix up to 200 kg N ha<sup>-1</sup> yr<sup>-1</sup> [2].

The elevated nutrient concentration in the leaves (especially N, K, and Ca) makes this tree component an important reserve of bioelements, although it represents only a small percentage of whole tree biomass (see *table IV*). In the living cells of the leaves, major amounts of nutrients are accumulated in relation to the processes of transpiration and photosynthesis [15].

Major concentrations of Ca are found in the bark (*table III*); Bodalla also has high concentrations of Ca in the leaves. According to Attiwill et al. [1], Ca plays a major role in the process of cell wall lignification and is not redistributed to new plant tissues. Various authors have proved that bark is the tree component with the highest concentrations of Ca. For example, Sharma and Pande [30] found this to be the case in hybrid *Eucalyptus* in 5 and 7 years old stands, and in *Acacia auriculiformis* in 3, 5, 7 and 9 years old stands; Campos [9] in *Ilex paraguariensis* (12 years); Leles et al. [16] in *Eucalyptus*

*camaldulensis* and *Eucalyptus pellita* (4.3 years); Pereira et al. [22] in *Acacia mearnsii* (9 years); and Vezzani [35] in *Acacia mearnsii* (3.7 years).

The highest concentrations of Mg were also found in leaves (*table III*), which have already been proved in several species at different stand ages [8, 10, 13, 22, 33]. According to Kramer and Kozłowski [15], leaf concentrations of Mg and Ca vary significantly with the season.

The highest concentrations of P and K are found in the leaves, whereas the lowest are in the dead branches probably due to a nutrient shift (resorption) into younger plant tissues or leaching during the dying process of the branches. However, the lowest concentrations of N, Ca, Mg, and S are found in the wood which implies that this is generally rich in C, H, and O (*table III*).

Mean nutrient contents in the above-ground biomass of the Bodalla are shown in *table IV*. Nutrient content in the total biomass components of the Bodalla follows the order: N > K > Ca > Mg > S > P.

**Table III.** Nutrient concentrations with their respective standard deviations ( $\pm$  s.d.) in tree components of Bodalla (AGROSETA Florestal Company, experimental sites, Rio Grande do Sul, Brazil).

Components	Nutrients (g kg <sup>-1</sup> )					
	N	P	K	Ca	Mg	S
leaves	23.6	1.00	9.20	6.80	1.60	1.18
( $\pm$ s.d.)	0.3	0.003	0.1	0.1	0.003	0.002
living branches	8.00	0.40	6.20	3.90	1.00	0.60
( $\pm$ s.d.)	0.2	0.002	0.2	0.008	0.003	0.002
dead branches	4.50	0.10	2.30	4.20	0.70	0.20
( $\pm$ s.d.)	0.01	0.0004	0.006	0.008	0.002	0.0005
bark	11.4	0.42	4.60	6.20	1.10	0.62
( $\pm$ s.d.)	0.2	0.0004	0.5	0.1	0.003	0.002
wood	2.30	0.14	3.30	0.70	0.20	0.12
( $\pm$ s.d.)	0.003	0.001	0.1	0.001	0.007	0.0004

**Table IV.** Mean nutrient contents in different components of above-ground biomass of Bodalla (AGROSETA Florestal Company, experimental sites, Rio Grande do Sul, Brazil). In parenthesis, percentage on total above ground biomass.

cp <sup>1</sup>	ag <sup>2</sup> biomass (kg ha <sup>-1</sup> )	Nutrients in the biomass (kg ha <sup>-1</sup> )					
		N	P	K	Ca	Mg	S
l <sup>3</sup>	4377 (22.5%)	103	4.38	40.4	29.8	7.2	5.2
lb <sup>4</sup>	3851 (19.8%)	30.7	1.54	24.0	15.2	4.0	2.3
db <sup>5</sup>	75.4 (0.4%)	0.3	0.007	0.2	0.3	0.1	0.02
b <sup>6</sup>	2421 (12.4%)	27.6	1.02	11.2	15.1	2.8	1.5
w <sup>7</sup>	8751 (44.9%)	20.3	1.22	28.6	6.3	2.1	1.1
t <sup>8</sup>	19475.4	181.9	8.17	104.4	66.7	16.2	10.1

<sup>1</sup> Components of above-ground biomass; <sup>2</sup> above-ground biomass of each component; <sup>3</sup> leaves; <sup>4</sup> living branches; <sup>5</sup> dead branches; <sup>6</sup> bark; <sup>7</sup> wood; <sup>8</sup> total above-ground biomass.

**Table V.** Percentage of accumulated nutrients in the components of above-ground biomass of Bodalla (AGROSETA Florestal Company, experimental sites, Rio Grande do Sul, Brazil).

Components	Nutrients (%)					
	N	P	K	Ca	Mg	S
leaves	56.8	53.0	39.0	45.0	45.0	51.0
living branches	17.0	19.0	23.8	23.0	24.0	23.0
dead branches	0.2	0.1	0.2	0.5	0.3	0.2
bark	15.0	12.9	11.0	22.5	17.0	12.0
wood	11.0	15.0	26.0	9.0	13.7	13.8

This result is similar to that found by Baggio and Carpanezzi [4] in *Mimosa scabrella*. Pereira et al. [22] found the following ranking of nutrients in above-ground biomass in a 9-year-old stand of *Acacia mearnsii*: N > Ca > K > Mg > P.

The different ranking of Ca in the latter study is due to an enrichment of Ca in the bark at the end of the forest rotation age (7 years).

In terms of nutrient exportation it is recommended that only trees (including bark) bigger than a diameter of 6.0 cm should be harvested. In this case, a considerable amount of residues remain on the forest floor as a protective layer between the atmosphere and the soil.

Caldeira [7] pointed out that there are no significant differences in nutrient export with respect to different harvest strategies of three Australian provenances (Bodalla, Lake George, and Batemans Bay) of *Acacia mearnsii*.

Baggio and Carpanezzi [4] analysed nutrient distribution in above-ground biomass of *Mimosa scabrella* in southern Brazil and found that 73% of all nutrients (N, P, K, Ca, Mg, S, Cu, Fe, Mn, and Zn) was found in the wood and only 27% in the crown. However, several authors have shown that tree crowns contain about 50% of the nutrients, with a major concentration in the leaves [22, 23, 27, 28, 30, 36].

In the present study, leaves account for more than 50% of the total content of N, P, and S (table V). It can also be illustrated that nutrient quantity in the leaves is superior to that of other above-ground biomass components, such as wood, although wood comprises the greatest proportion of the above-ground biomass [7]. This is usual in boreal forests but can also be seen in temperate forests [14, 37].

Considering the usual subdivision into crown and trunk biomass, 43% of the dry matter was composed of leaves and living and dead branches, with the following above-ground biomass contents: 74% of N, 73% of P, 62% of K, 68% of Ca, 70% of Mg, and 75% of S. The trunk components (bark and wood) accumulated 26% of N, 27% of P, 38% of K, 32% of Ca, 30% of Mg, and 25% of S (table V). In terms of nutrient distribution Vezzani [35] obtained similar results in another study in a 3.7-year-old stand of *Acacia mearnsii*.

Nutrient measurements are important in the analysis of ecologically sustainable management systems of tropical and subtropical plantations; e.g., harvest strategies can be optimised by taking into account the different nutrient contents of tree components in different stand stages. In particular, harvests in very young stands (so-called “mini-rotations”) with high amounts of juvenile wood can lead to higher nutrient export compared to older stands.

#### 4. CONCLUSIONS

According to the previously described results on *Acacia mearnsii*, provenance Bodalla, it can be concluded that:

- the highest nutrient concentrations were found in the leaves;
- the lowest concentrations of P and K were found in the dead branches, and those of N, Ca, Mg, and S in the wood;
- the following nutrient distribution was found in the trunk biomass (bark and wood): 26% of N, 27% of P, 38% of K, 32% of Ca, 30% of Mg, and 25% of S. By reverse, 74% of N, 73% of P, 62% of K, 68% of Ca, 70% of Mg, and 75% of S of the total above-ground biomass were found in the leaves, living and dead branches (mostly crown).

This work is a first approach towards optimising management in South Brazilian plantations of *Acacia mearnsii*; further investigations are necessary, especially relating to crop rotations, below-ground biomass, and nutrient dynamics, such as fluxes and resorption processes.

**Acknowledgements:** We thank Mr. Elias Moreira dos Santos from AGROSETA S.A. and the staff of the Forestry Department of the University of Santa Maria for their technical support. Furthermore, we thank the reviewers for very helpful comments on the manuscript.



## REFERENCES

- [1] Attiwill P.M., Guthrie H.B., Leuning R., Nutrient cycling in a *Eucalyptus obliqua* (L'Herit) forest. I. Litter production and nutrient return, *Aust. J. Bot.*, Melbourne 261 (1978) 79–91.
- [2] Auer C.G., Silva R., Fixação de nitrogênio em espécies arbóreas, in: Cardoso E.J.B.N., Tsai S.M., Neves M.C.P. (Eds.), *Microbiologia do Solo*, Campinas, Sociedade Brasileira de Ciência do Solo, 1992, pp. 157–172.
- [3] Baggio A.J., Estudio sobre el sistema agroforestal tradicional de la bracinga (*Mimosa scabrella* Benth.) en Brasil: Productividad, manejo de residuos y elaboración de compost, Madrid, Doctoral Thesis, Universidad Politécnica de Madrid, 1994, 242 p.
- [4] Baggio A.J., Carpanezzi A.A., Estoque de nutrientes nos resíduos da exploração de bracingais, *Boletim de Pesquisa Florestal*, Colombo 34 (1997) 17–29.
- [5] Bellote A.F.J., Suprimento de nutrientes minerais e crescimento de plantações adubadas de *Eucalyptus grandis* nos cerrados do Estado de São Paulo, Freiburg, Doctoral Thesis, Universität Freiburg, 1990, 166 p.
- [6] Bellote A.F.J., Sarruge J.R., Haag P.H., Oliveira G.D., Extração e exportação de nutrientes pelo *Eucalyptus grandis* Hill ex Maiden em função da idade. 1 – Macronutrientes. IPEF, Piracicaba, 20 (1980) 1–23.
- [7] Caldeira M.V.W., Quantificação da biomassa e do conteúdo de nutrientes em diferentes procedências de Acácia-negra (*Acacia mearnsii* De Wild.), Santa Maria-RS, Master Thesis, Universidade Federal de Santa Maria, 1998, 120 p.
- [8] Caldeira M.V.W., Pereira J.C., Schumacher M.V., Della-Flora J.B., Santos E.M., Comparação entre as concentrações de nutrientes nas folhas e no folheto em procedência de *Acacia mearnsii* De Wild. *Revista Árvore*, Viçosa 23 (1999) 489–492.
- [9] Campos M.A.A., Balanço de biomassa e nutrientes em povoamentos de *Ilex paraguariensis*. avaliação na safra e na safrinha. Curitiba-PR, Master Thesis, Universidade Federal do Paraná, 1991, 106 p.
- [10] Drummond M.A., Barros N.F., Souza A.L., Silva A.F., Distribuição de biomassa e de nutrientes em diferentes coberturas florestais e pastagem na região do médio Rio Doce-MG. *Revista Árvore*, Viçosa 21 (1997) 187–199.
- [11] Empresa Brasileira de Pesquisa Agropecuária (EMBRAPA). Levantamento de reconhecimento dos solos do Estado do Rio Grande do Sul. EMBRAPA, Recife (Boletim Técnico, 30), 1973.
- [12] George M., Varghese G., Nutrient cycling in *Eucalyptus globulus* plantations: I. Organic matter production, nutrients accumulation in standing crop and nutrients removal through harvest, *Indian For.* 116 (1990) 42–48.
- [13] Gomes F.S., Pessotti J.E.S., Pacheco R.M., Exportação de nutrientes por clones de *Eucalyptus urophylla*, em três unidades de solo no Vale do Rio Jarí, in: IUFRO Conference on Silviculture and Improvement of Eucalypt, Salvador, BA, Proceedings, v.3, 1997, pp. 209–214.
- [14] Kimmins J.P., *Forest Ecology*, Collier Macmillan Canada, New York, 1987, 531 p.
- [15] Kramer P.J., Kozlowski T., *Physiology of woody plants*, Academic Press New York, 1979, 811 p.
- [16] Leles P.S.S., Reis G.G., Reis M.G.F., Distribuição de nutrientes em *Eucalyptus camaldulensis* e *Eucalyptus pellita* sob diferentes espaçamentos, na região de cerrado, in: Congresso Brasileiro de Ciência do Solo, Viçosa, MG, Resumos expandidos, Viçosa, 1995, pp. 862–863.
- [17] Lugo A.E., Comparison of tropical tree plantations with secondary forests of similar age, *Ecol. Monogr.* 62 (1992) 1–41.
- [18] Miller H.G., Internal and External Cycling of Nutrients in Forest Stands, in: Pereira J.S., Landsberg J.J. (Eds.), *Biomass Production by Fast-Growing Trees*, Applied Science Vol. 166, Kluwer Academic Publ., Dordrecht, 1989, pp. 73–80.
- [19] Moreno J.A., *Clima do Rio Grande do Sul*, Secretaria da Agricultura, Porto Alegre, 1961.
- [20] Nambiar E.K.S., Brown A.G. (Eds.), *Management of Soil, Water and Nutrients in Tropical Plantation Forests*. Australian Center for International Agricultural Research (ACIAR), Monograph No. 43, Canberra, 1997, 571 p.
- [21] Oliveira J.B., Jacomine P.K., Camargo M.N., Classes gerais de solos do Brasil, Jaboticabal, FUNEP, 1992, 201 p.
- [22] Pereira J.C., Caldeira M.V.W., Schumacher M.V., Hoppe J.M., Santos E.M., Determinação do conteúdo de nutrientes em um povoamento de *Acacia mearnsii* De Wild. no Rio Grande do Sul. *Revista Árvore*, Viçosa 24 (2000) 193–199.
- [23] Pereira J.C., Schumacher M.V., Hoppe J.M., Caldeira M.V.W., Santos E.M., Produção de biomassa em um povoamento de *Acacia mearnsii* De Wild. no Estado do Rio Grande do Sul, *Revista Árvore*, Viçosa 21 (1997) 521–526.
- [24] Poggiani F., Ciclagem de nutrientes em ecossistemas de plantações florestais de *Eucalyptus* e *Pinus*. Implicações silviculturais, Tese de livre-docência, ESALQ-USP, 1985, 211 p.
- [25] Poggiani F., Alterações dos ciclos biogeoquímicos em florestas, in: Congresso Nacional sobre Essências Nativas, 1992, São Paulo, SP, Proceedings, 1992, pp. 734–739.
- [26] Poggiani F., Couto H.T.Z., Suiter Filho W., Biomass and Nutrient Estimates, in: Short Rotation Intensively Cultured Plantation of *Eucalyptus grandis*, Piracicaba, IPEF 23 (1983) 37–42.
- [27] Reis M.G.F., Barros N.F., Ciclagem de nutrientes em plantios de eucalipto, in: Barros N.F., Novais R.F. de (Eds.), *Relação solo-eucalipto*, Viçosa, Ed. Folha de Viçosa, 1990, pp. 256–296.
- [28] Rezende G.C., Barros N.F., Moraes T.S.A., Mendes C.J., Produção e macronutrientes em florestas de eucalipto sob duas densidades de plantio, *Revista Árvore*, Viçosa 7 (1983) 165–196.
- [29] Schumacher M.V., Ciclagem de nutrientes como base da produção sustentada em ecossistemas florestais, in: Simpósio sobre Ecossistemas Naturais do Mercosul: O Ambiente da Floresta, Santa Maria, RS, Proceedings, 1996, pp. 65–77.
- [30] Sharma S.C., Pande P.K., Patterns of litter nutrient concentrations in some plantation ecosystems, *For. Ecol. Manage.* 29 (1989) 151–163.
- [31] Tandon V.N., Pande M.C., Singh R., Biomass estimation and distribution of nutrient in five different aged *Eucalyptus grandis* plantations in Kerala state, *Indian For.* 114 (1988) 184–199.
- [32] Tedesco M.J., Volkweiss S.J., Bohnen H., Análise de solos, plantas e outros materiais, Porto Alegre: UFRG, Departamento de Solos/Faculdade de Agronomia (Boletim Técnico, 5), 1985, 188 p.
- [33] Turvey N.D., Smethurst P.J., Nutrient concentrations in foliage, litter and soil in relations to wood production of 7 to 15 year old *Pinus radiata* in Victoria, Australia, *Aust. For.*, Canberra, 57 (1994) 157–164.
- [34] Van den Driessche R., Prediction of mineral status of trees by foliar analysis, *Bot. Rev.*, New York, 40 (1984) 347–394.
- [35] Vezzani F.M., Aspectos nutricionais de povoamentos puros e mistos de *Eucalyptus saligna* (Smith) e *Acacia mearnsii* (De Wild.), Porto Alegre-RS, Master Thesis, Universidade Federal do Rio Grande do Sul, 1997, 97 p.
- [36] Waring R.H., Schlesinger W.H., *Forest ecosystem, concepts and management*, Academic Press, Orlando, 1985, 340 p.
- [37] Whittaker R.H., Borman F.H., Likens G., Siccama T.G., The Hubbard Brook ecosystem study: forest biomass and production, *Ecol. Monogr.* 44 (1979) 233–252.
- [38] Young H.E., Carpenter P.N., Sampling variation of nutrient element content within and between on trees of the same species, in: *Oslo Biomass Studies*, Oslo, 1976, pp. 75–90.